

UNSURE ABOUT CASTRATION?

Suprelorin® 4.7 mg is suited to dog owners who want a temporary non-surgical solution for castration.

BENEFITS OF CHEMICAL NEUTERING

- ✓ Temporary reproductive control.
- ✓ May alleviate diseases that are dependent on testosterone (e.g. prostatic enlargement).
- ✓ May address testosterone related behavioural problems.


Behaviour problems are not always due to testosterone. Please talk to your vet who will be able to advise if this product is best for your dog.

MODE OF ACTION

Pituitary Gland

Drug acts here: down-regulation of GnRH receptors

Implant inserted here



DID YOU KNOW:

- ✓ Implantation does not require anaesthetic.
- ✓ The implant does not have to be removed.
- ✓ One 4.7 mg implant results in at least 6 months of infertility in male dogs.
- ✓ Male dogs return to complete fertility once the active has worn off.

DO YOU HAVE QUESTIONS? PLEASE ASK YOUR VET TODAY.

Visit nz.virbac.com/suprelorin
for more information

Shaping the future
of animal health



Suprelorin® 4.7 mg is a Restricted Veterinary Medicine. Available only under Veterinary Authorisation. Registered Pursuant to the ACVM Act 1997, No. A009158. Copyright © 2022 Virbac New Zealand Limited. All rights reserved. Virbac New Zealand Limited, 26-30 Maui Street, Pukeke, Hamilton, 3200. 5PR0025. 08/22.

Suprelorin® 4.7 mg

KEEP THEIR OPTIONS OPEN



The temporary contraceptive
implant for male dogs.

Shaping the future
of animal health



WHAT IS SUPRELORIN® 4.7 mg?

Suprelorin® 4.7 mg is a temporary contraceptive implant for male dogs, administered by your veterinarian.

It is inserted under the skin between your dog's shoulders.

It prevents testosterone and sperm production.

There is no need for anaesthetic or surgery.

HOW DOES IT WORK?

After implant placement, an active ingredient is steadily released preventing the production of testosterone.

Testosterone is essential for sperm production and reproduction in the male.

It takes around 6 weeks for the dog to become completely infertile due to remnant sperm stored in the reproductive tract.

Once infertile, sperm production is prevented for at least 6 months. Once the implant wears off the dog's fertility will gradually return.*

What will you see?

- Loss of interest in bitches.
- Loss of ability to ejaculate.
- Smaller testicles.

These are all temporary and will all return to normal as the active wears off.

Safety

Suprelorin® 4.7 mg is safe and not associated with significant side effects. To ensure continued infertility it is recommended to place an implant every 6 months.

*Discuss with your veterinarian prior to use in a breeding dog. In some dogs infertility can last longer than 6 months.

WHY USE SUPRELORIN® 4.7 mg?

- ✓ To trial the effects of neutering prior to permanent surgery.
- ✓ Test the behavioural response to lowered testosterone.
- ✓ For the benefit of reproduction control now, with the option to breed in future*.
- ✓ To still have the benefits of neutering when anaesthesia and surgery is not ideal.

